

List of figures

- I. Staphylococcal strain preserved in nutrient agar slant.**
- II. Catalase test showing reaction.**
- III. Mannitol Salt Agar Media of inoculation the bacterial strain.**
- IV. Antibiotic sensitivity test of the bacterial strain on Muller-Hinton agar.**
- V. Slide coagulation test of the bacterial strain performed against human blood.**
- VI. Gram staining of the bacterial strain seen under phase contrast microscope.**
- VII. Bioassay of the Crude protein with human blood samples obtained by acetone precipitation.**
- VIII. Bioassay of the Crude protein with human blood samples obtained by ammonium sulphate precipitation.**
- IX. Slide Coagulation test of the lyophilised fraction of the coagulase protein extracted from the bacterial strain performed against human whole blood.**
- X. Changes in the prothrombin time after the treatment with the lyophilised protein.**
- XI. Changes in the thrombin time after the treatment with the lyophilised protein.**
- XII. Changes in the activated partial thromboplastin time after the treatment with the lyophilised protein.**
- XIII. Changes in the fibrinogen time after the treatment with the lyophilised protein.**
- XIV. Stability of coagulase protein at different temperature.**
- XV. SDS PAGE of Staphylocoagulase Protein.**
- XVI. Standard graph of individual amino acids.**
- XVII. Chromatogram showing amino acids of coagulase protein fraction having molecular weights of 66 kDa using HPLC.**
- XVIII. Chromatogram showing amino acids of coagulase protein fraction having molecular weights of 33 kDa using HPLC.**

- XIX. Chromatogram showing amino acids of coagulase protein fraction having molecular weights of 14 kDa using HPLC.**
- XX. Protein model generated through MODELER 9v8 (MODELS view through Discovery Studio 4.0 molecule viewer).**
- XXI. Ramachandran plot of the generated model of the coagulase protein.**
- XXII. Disorder Prediction through “DISOPRED” for protein disordered region detection.**
- XXIII. Results of ToxinPrediction for Toxicity of the coagulase protein.**
- XXIV. Active sites of generated protein model with residues view through Discovery Studio 4.0**
- XXV. The PCR amplified product in agarose gel (1%) using bacteria specific primers.**
- XXVI. Phylogenetic tree of *Staphylococcus NEIST basab -1(S 43)* in MEGA 3.1 software using neighbour joining method**