

CHAPTER-X

BIBLIOGRAPHY

BIBLIOGRAPHY

References :

- Ali, S.(1982) Effect of hatching of eggs of Bombyx locust, *Patang succinita*, (Orthoptera: Acrididae), Comp. Physiol. Ecol., 7(2):78-80.
- Annual Report, CSR & TI, Berhampore,(2000-2001)
- Arunachalam, V. and Bandopadhyay, A. (1984) A method to make decisions jointly on a number of dependent characters. *Indian J. Genet.*, 44 (3):419-424.
- Begum Naseema A.; Joge, P.G., Basavaraja, H.K., Karippa, B.K., Nanje, Gowda, B.,and Dandin, S.B. (2003) Evaluationn of polyvoltine x Bivoltine Hybrids of Silkworm, *Bombyx mori* L. Proceedings of the National Conference on Tropical Sericulture for Global Competitiveness :105-112.
- Bencharmin,K.V.;Manjeet Jolly, S., Benjamin, D.A.I. (1988). Studies on the reciprocal crosses of multivoltine and bivoltine breeds in silkworm *Bombyx mori* L. with special reference to the use of bivoltine hybrid as a parent. *Indian Journal of Sericulture*, 27(1):27-34.
- Bhattapadhyay, G. K.; Sengupta, A.K., Verma, A. K., Sen, S. K. and Charatchandra,B.(2001b) Transgression of shell weight- A multigenic trait, through development of congenic breed in tropical silkworm *Bombyx mori* l. *Sericologia*.41(1):33-42.
- Chakravorty , R.(2002) Studies on the adaptability of some improved mulberry varieties in the Agroclimatic condition of Assam with special reference to leaf yield and their utility for silkworm rearing. Ph.D. Thesis, Gauhati University, Guwahati.
- Chang. K.Y.; Han.K.S.and Min.B.Y.(1981) Genetic studies on silkworm characters by diallel cross II. Analysis of heterosis and combining abilities. *Seric.J.Korea*,22(1): 68.
- Chatterjee,S.N.(1992a)Silkworm rearing Brain Storming Meeting on Genetics and Bio-technology in Silkworm and Mulberry, CSR & TI, Mysore, May 28-29.

Chatterjee, S. N.,(1992b)Silkworm improvement and rearing. Natl. Conf. Mulb. Seri. Res., CSR & TI, Mysore, December, 10-11.

Chatterjee, S.N.and Datta R.K. (1992). Hierarchical clustering of 54 races and strain of mulberry silkworm, *Bombyx mori* significance of biochemical parameters. Theor. Applications Genet. 85 : 394-402.

Chatterjee, S.N.; Rao, P.R. M; Jayaswal, K.P.; Singh Ravindra and Datta,R.K.(1993) Genetic variability in mulberry silkworm *Bombyx mori* L.breeds with low silk yield. *Indian J.Seric.*,**32**:69- 86

Chowdhury, S. N.(1992) *Silk and Sericulture*, Govt. of Assam, 17-22.

Congdon,B.D. and Logan, J. A. (1983) Temperature effect on development and fecundity of *Oligonychus pratensis*, *Envr.Ent.*, **12**:359-362.

Cunnington, A.M. (1985) Factors affecting oviposition and fecundity in the grain mite, *Acarus siro* (Acarina Acaridae) especially temperature and humidity. *Exp.App. Acarol.l.(4)*: 327-344.

Das,S.K, Pattnaik, S., Ghosh, b., Singh, T., Nair, B.P., Subba Rao,G.(1994) Heterosis analysis in some three way crosses of *Bombyx mori* LSericologia,**34**(1):51-56.

Dash, A. K., Patro,P.C., Nayak,B.K.and Dash, M.C. (1994) Cocoon crop performance of the Indian wild tasar silkworm *A.papahia* (*Lepidoptera:Saturniidae*) reared on different food plants. *Int.J.Wild Silkmoth and Silk***1**:72-74.

Datta,R.K.,Sureshkumar,N,Basavaraja,H.K.,Mal Reddy,N.(2000). "CSR18 X CSR19"-a robust bivoltine hybrid for rearing throughout year in tropics. Abstract in seminar on sericultural technology an appraisal held at CSR&TI, Mysore, and Karnataka:**6-7**.

Datta , R.K. (2000) Silkworm breeding in India : Present status and new challenges . Paper presented in National Conference on strategies for Sericulture Research and Development, CSR&TI , Mysore, India,November 16-18

Datta, R.K.; Basavaraja, H.K., Mal Reddy, N., Nirmal Kumar, S., Suresh Kumar, N., Ramesh Babu, M., Ahsan M, M. and Jayaswal, K. P. (2001) Breeding of new productive bivoltine hybrid, CSR 12 CSR 6 of silkworm *Bombyx mori* L. *Int. J. Indust. Entomol.* 3 : 127-133.

Devaiah, M.C.; Narayanaswamy, K.C. and Maribashetty, V.G. (1999) *Advances in mulberry sericulture*, Bangalore, India, 190-193, 525-528.

Dorcas, D.and Vivekanandan, M. (1991) Screening of mulberry varieties for rainfed condition. *Sericologia*,**31(1)**:223-241

Falconer , D.S. (1952) The problem of environment and selection. *Am.Nat.*, **86**:293-309.

Falconer , D.S. (1960) Selection mice for growth on high and low planes of nutrition. *Genet. Res.*,**1**:91-113.

Falconer , D.S. (1990) Selection in different environments : effects on environmental sensitivity (reaction norm) and on mean performance. *Genet. Res.*,**56**:57-70i.

Gamo, T. and Ohtsuka, Y. (1980) Phylogenetic studies on the racial differences of the silkworm, *Bombyx mori* on the basis of polymorphic genes in haemolymph proteins. *Bull. Sericult. Exp. Sta.*, **28**:15-50.

Gamo, T. (1983) Genetic analysis of growth rate, pupation rate and some quantitative characters by diallele analysis in the silkworm. *Jpn. J. Breed.*, **33(2)**:178-190

Gamo, T.and Hirabayashi, T. (1983)Genetic analysis of growth rate, pupation rate and some quantitative characters by diallele cross in the silkworm *Bombyx mori* L. *Japan.J.Breed.*, **33**:179-180

Ghosh, S.P.(1991) *Agro-climatic Zone efic Research Indian Perspective under NARP*.Indian Council of Agricultural Research, New Delhi,539.

Hajare, T. N.; Prasad, J., Jadav, A. D. and Challa, O. (2007) Influence of soils on productivity of sericulture in region of Maharashtra, India. *Soils and Crops*, 17 (1):186-187.

Hajare,T.N.; Jadav,A.D., Jadish Prasad, Patil,N.G.and Lal Sohan. (2008) Performance of silkworm breeds, *Bombyx mori L. Indian J. Seric.*, 47:111-114.

Harda, C. (1949) On the hybrids effects of economical characters in silkworm, *J.Seric.Exp.Stn.*,17(1):50-52.

Harada C (1961) Heterosis of quantitative characters in the silkworm. Bull. Sric. Expl. Stan., 17:1-52.

Hee, S.M., Yan . X., Mi , Y., Xia, L .(1989) Breeding of a new silkworm variety “ Feng1 x 54a” for summer-autumn rearing *Sci Seric.*, 15, 79-87.

Hodai, T. (1949) Effect of environmental humidity on embryonic development and hatching in the eri silkworm with consideration on the mechanism, *J. Sericult. Sci.Japan*,18,171-180.

Jain, A. K.; Sarkar, A. and Bajpai, A. K. (1992) Screening of tolerance to salinity and alkalinity in mulberry through merited culture. *National Conf. Mulberry Seric. Res.*,32-35.

Johansen, N.B.(1997) Influence of temperature development, fecundity and survival of cabbage moth *Mamestra brassicae* (L) (Lepidoptera: Noctuidae) in relation to the improvement of forecasting and control methods, *J. Appl Ent.*,121,81-88.

Jolly, M.S., Chaturvedi, S. N. and Prasad, S. (1968)A Survey of Tassar Crops in India.*Ieri*.7(1):56-65.,*CSR & TI, Mysore, Dec10-11*,:14

Kempthrome, O. (1952). *The Design and Analysis of Experiments*. New York : Wiley.

Khan, M. A. (2006) Mulberry silkworm crop production management and improvement in sericulture as an economic enterprise in North West India” held at RSRS, Dehradun on Nov.11:159-168.

Khanna, S.S., (1989)The agro-climatic approach. In : Survey of Indian agriculture, The Hindu, Madras, India, 28-35.

King, A.B.S. (1975) Factors the first brood of affecting the phenology of the sugarcane froghopper *Aeneolamia varia saccharina* (Homoptera: Cercopidae) in Trinidad. Bull.Ent. Res.**65**:359-372.

Kiran Kumar K. P. and Sankar Naik,S. (2011) Development of polyvoltine x bivoltine hybrids of mulberry silkworm, *Bombyx mori* L. Tolerant to BmNPV. International Journal of Zoological Research. 7:300-309.

Kiran Kumar K. P., Sankar Naik S. (2011) Development of polyvoltine x bivoltine hybrids of mulberry silkworm, *Bombyx mori* L. Tolerant to BmNPV. International Journal of Zoological Research. 7:300-309.

Krishnaswami, S.; Asan, M. and M. and Sriharan, T.P.(1970) Studies on the quality of mulberry leaves and silkworm cocoon crop production. *Indian J. Sericulture*, **9(1)**: 11-25.

Krishnaswami, S.;Sriharan, T. P. and Ahasen, A.M.(1971) Ecological studies on silkworm rearing to prevent crop losses in adverse seasons in West Bengal, *Indian J.Seri.*,**10(1)**:72-76.

Krishnaswami, S. and Narasimhanna, M . N . (1974) Large-scale trials on bivoltine hybrids in Mysore state. *Indian. J Genetics*, **34(A)** : 229-236.

Krishnaswami, S. (1978) New Technology of Silkworm Rearing, CSR & TI, Bulletin No.2, Central Silk Board, Bangalore, India,1-23

Krishnaswami, S. (1986) Improved Method of Rearing Young Age (chawki) Silkworms. Central Silk Board , Bulletin No.3,CSR&TI , Mysore.

Krishna Rao , S (1998) Bivoltine silkworm races evolved by KSSRDI. In "Silkworm Breeding". (Eds) G . Sreramareddy Oxford & IBH Pub. Co. Pvt. Ltd.

Krishna Rao, S.; Rajendra undukar, Raghu Raman, R., and Bongale, U. D (2006)Breeding strategies for robustness. In proceedings: Silkworm Breeders meet at CSR&TI, Berhampore,February 14-15, 3-20.

Kumar Suresh N.;Palit,A.K.;Kalpana ,G.V.and Joge,P.G.(2003) New index method for evaluation of silkworm hybrids of *Bombyx mori* L. Advances in mulberry sericulture,74-78.

Kumar Kishor;C.M. Basavaraja, Dutta, R.K., Raghavendra Rao, D.,Jayaswal , K.P., Premalatha , Singh , Ravindra and kariappa, B.K. (2001) Heterosis inrelation

to combining ability in multivoltine and bivoltine strains of silkworm, *Bombyx mori* L. *Indian J.Seric.*,**40**: 1-6.

Kumar Rajesh and Elangovan Vadmalai (2013) Rearing performance of Eri Silkworm (*Philosamia ricini*) during Autumn season of Uttar Pradesh. *Indian J.Seri.*,**52(2)**:131-138.

Kumaresan, P.; Sinha, R. K., Sahni, N.K. and Seka, S.(2000)Genetic variability and selection indices for economic quantitative traits of multivoltine mulberry silkworm (*Bombyx mori* L.) genotypes, *Sericologia*, **40**: 595-605.

Mano, Y.; Nirmal Kumar, S. Basavaraja , H.K. Mal Reddy, N and Dutta, R.K. (1992) An index for multiple trait selection or silk yield improvement in *Bombyx mori* L. *Natl. Conf. Mulb. Seric. Res.*, CSR & TI, Mysore,116.

Mano; Nirmal Kumar, S., Basavaraj, H. K., Mal Reddy, N., Datta, R. K. (1993) Anew method to select promising silkworm breeds/combinations. *Indian Silk*,53.

Mano; Y., Nirmal Kumar, S., Basavaraja, H.K.,MalReddy, N. and Datta, R.K. (1997) Combined trait selection index for silk yield improvements in *Bombyx mori* L. In : Silkworm Breeding, Ed.G.Sreerama Reddy ,Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi, 278-284.

Mathur, S.K.; Parmanik, D.R., Sen, S .K. and Subba Rao, G (1975) Effect of seasonal temperature and humidity on ovulation, fecundity and retention of eggs in silkmoth *Bombyx mori* (L) (Lepidoptera: Bombycidae) *Rec. Zool. Surv. India*, **95 (1- 2)**:57-64.

N.Suresh Kumar; H. K. Basavaraja, P. G. Joge, N. Mal Reddy, G. V. Kalpana and S.B. Dandin. (2006) Development of a new robust bivoltine hybrid (CSR46 x CSR47) of *Bombyx mori* L. for the Tropics.

Nagaraju, J.; Premalatha, V. Sing, Ravindra, Noamani, M. K. R .and Jolly, M. S. (1989). Isolation of a polyvoltine strain with sex-limited larval marking in silkworm *Bombyx mori* (Lepidoptera: Bombycidae. *Sericologia*, **29**:395-502.

Nagaraju, J.; Raje Urs, S.and Dutta, R.K.(1996) Cross-breeding and heterosis in silkworm, *Bombyx mori* L. A review. *Sericologia*, **36**;1-26 Sing, Ravindra; Nagaraju,J.; Ram Mohan Rao P.; Premalatha,V.; Vijayaraghavan, K. and Sengupta, K. (1990). Heterosis analysis in the silkworm, *Bombyx mori* L. *Sericologia*, **30**:293-300.

News & Views, a quarterly bulletin of CSR&TI, Berhampore.(1989) Vol.4 (2).

Nirmal Kumar, S.; Ramesh Babu, M., Basavaraja, H. K., Mal Reddy, N. and Datta, R. K. (1998) Double hybrid for improvement of silk production in silkworm *Bombyx mori* L. In: *Silkworm Breeding*, G. Sriramreddy (Ed.), Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, 209-217.

Nirupama , R. and Ravindra Singh. (2007) Evaluation of polyvoltine breeds of mulberry silkworm *Bombyx mori* L. *J. Exp. Zool, India* **10(2)**: 341-344.

Quadir, S.M.; Mir Nisar, Khan, M.A. and Ahsan, M.M. (1997), Identification of season specific silkworm hybrids for temperate climatic condition of Kashmir, Abstt. No. P-2, National conference on strategies for sericulture research and Development, CSR&TI., Mysore, India, 16-18.

Raghavendra Rao , D.;Premalatha , V.Singh Ravindra ., Kariappa, B. K., Jayaswal, K.P., Ram Mohana Rao, P. and data, R.K. (2001) Studies on manifestation of hybrid vigour in polyvoltine x bivoltine crosses of silkworm, *Bombyx mori* L. *J. Exp. Zool.*, 4:219-232.

Raghavendra Rao, D.; Singh, Ravindra., Kariappa, B.K., Premalatha, V. (2002) Studies on manifestation of hybrid vigour and cocoon shape variation in multivoltine x bivoltine hybrids of the silkworm, *Bombyx mori* L.*Uttar Pradesh J.Zool.*,22:141 -147.

Rahman,M.;Khalequzzaman and Rahman, S.M.(1980) Studies on the oviposition, hatchability by some multivoltine race of the silkworm *Bombyx mori* L.Lepidoptera: Bombycidae), *Indian J.Seri.*,**19 (1)**:28-30.

Rajadurai,S.;Philip, Tomy and Shekhar, M.A.(2010) Seasonal rearing performance of Eri silkworm *Samia Cynthia ricini* (B) on castor and Tapioca under South Karnataka condition. *Indian J.Seric.*,**49 (2)**:134-137.31.

Raju. P.J.; Guruswamy,D., Raghuraman ,R., and Bongale,U.D.(2003) New productive multivoltine hybrids of silkworm *Bombyx mori* for commercial exploitation under rainfed sericulture. Proceedings of the National Conference on Tropical Sericulture for Global Competitiveness,November 5 -7,54-57.

Rama Mohana Rao, P.; Somashekhar, T.H., Ahsan, M. M. and Datta, R. K. (1995) Studies on the size of deviation in single cocoon filament in bivoltine breeds of the silkworm *Bombyx mori* L. Current Technology Seminar, September 20- 22, Central Sericultural Research and Training Institute, Mysore, India, 11.

Rama Mohana Rao, P.; Quadri, S. M. H., Nirmal Kumar, S., Premalatha, V., Dayananda, Joge P.G., Radhalakshmi, Y. C. P.G., Radhalakshmi, Y. C. and Shivakumar, K. P. (2012) Identification of polyvoltine x bivoltine hybrids of the *Bombyx mori* L. with superior fibre quality: a breakthrough in silkworm breeding.

Ramesh Babu, M.; Chandrashekaraiah, Lakshmi, H. and Prasad, J. (2001) Silkworm (*Bombyx mori*) genetic stocks-An evaluatory analysis. *Bull. Indian Acad. Seric.*, **5**:9-17.

Ranga Rao, G. V., Weightman, J. A. and Ranga Rao, D. V. (1989) Threshold temperatures and thermal requirements for the development of *Spodopter litura* (Lepidoptera : Noctuidae), *Environ. Entomol.* **18**:546-551.

Rao, P. R. M.; Singh, Ravindra., Jayaswal, K. P., Chatterjee, S. N. and Datta, R. K. (1998) Evaluation of some Indian and exotic low yielding silkworm (*Bombyx mori* L.) breeds through diallel cross and its significance for sericulture in dry zones. *J. Ent. Res.*, **22**:23-33.

Ravindra Singh ; Rachna Goel , Rao , D.R., Rekha, M., Premalatha, V., Ahsan, M.M. and Dutta , R.K. (2001) Evaluation of multivoltine x bivoltine F1 hybrids of the mulberry silkworm *Bombyx mori* L Sericologia, **41**: 231-238.

Regniere, J. and Turgeon, J. (1989) Temperature dependent development of *Zeiraphera canadensis* and stimulation of phenology. *Entomol. Exp. Appl.*, **50**: 185-193. Sinha, A.K.; U. S. P., Brahmachari, B. N., Ghosh, J. K. and Sinha, S. S. (1992) Race differences in cocoon shells of mulberry silkworm, *Indian J. Seric.*, **31(2)**: 167-168.

Roy, G. C.; Ghosh, B., Das, S. K., Nair, B. P., Rao, P. R. T., Sen, S. K., Sengupta, K. (1997) Comparative performance of multi x bi and six multi hybrids in commercial seasons of Eastern India. *Sericologia*, **37(1)**:1-18.

S. Kishore; N. Sivarami Reddy, S. Vidyunmala and N. Surryanarayanan. (2000) Evaluation of new mulberry hybrids under semi-arid condition of Andhra Pradesh, Abstt. No.P-30, National conference on strategies for sericulture Research and Development, CSR&TI., Mysore, India, 16-18, November, 2000.

Sekharappa, B.M.; P.G., Keshavareddy, K. S. and Dandin, S. B. (1999) Synthesis of CN2-A variable bivoltine silkworm race suitable for traditional rainfed area of Karnataka. Proc. XVIII, *Int.Serc.Cong.*, Oct12-16, Cario, Egypt.,97-104.

Sen, S. K.; Das, S. K., Rao, P.R.T., Ghosh, B., Das, N.K., Chattopadhy, S., Roy, G. C. and Sinha, S. S.(1995) on some important genetic parameters in the silkworm (*Bombyx mori* L.) *Indian J.Genet. Plant Breed.*,**55**:238-242.

Senapati, M. D. and Hazarika, A.(2012) Identification of some productive bivoltine hybrids of mulberry silkworm, *Bombay more*, L. through rearing under rainfed sericulture of North eastern region of India. *The Clarion.*, **1(1)**:93-104.

Senapati, M.D. and Hazarika, A. (2012) Studies of some economic traits of multi x bivoltine hybrids of mulberry silkworm, *Bombay mori* ,L. through rearing under rainfed condition for commercial exploitation in the agro climatic condition of Assam, India. *The Clarion.*, **1(2)**:142-147.

Senapati, M.D. and Hazarika, A.(2014) Evaluation of some productive hybrids of *Bombay more* ,L. suitable for the autumn seasons in the rainfed condition of Assam, India. *The Clarion.*,**3 (2)**:43-52.

Sengupta, K. (1968) An analysis of genotype environment interaction in same races of Silkworm, *Bomrax mori* L. *Indian J. Seric* **8 (1)** : 4-6.

Sharan, S. K.; M. K. Ojha., N. G. and Sinha, S. (1994) Regulation of voltinism in *A. mylitta* D. by manipulation of rearing period of larval stages.*Int.J.Wild Silkmoth & Silk* **1**:191-194.

Shao , Y .; H., Li, W.B., Xia , J-Q ., Cao , J.R.(1987) The breeding of the mulberry varieties “ Xinhang and Keming” for summer –autumn rearing. *Canye Kexua*, **13**: 15-21.

Shekar, Chandra; K. B. and Basavaraja H.K.(2011) Hybrid Vigour in Bivoltine Silkworm, *Bombyx mori* L., under different selection methods. *Indian J.Seri.*, **50(1)** : 28-31.

Siddiqui, A. A.; Debnath, A. K. and Sengupta, K. (1983) A study of genetic variability of some economic traits in *A. mylitta* D. Proc. National Seminar on Silk Research & Development, 41.

Siddiqui, A.; Sengupta,A. K., Kumar Ajit and, K. (1988) Genotypic and phenotypic variability of some quantitative characters in *A. mylitta*. *Sericologia*,**28(2)**:187-192.

Siddiqui, A.; Sengupta, A. K. (1994a) Genetic variability in technological characters-First report A.*mylitta*,*Sericologia*,**34(1)**:149-154.

Singh, Ravindra; Kalpana, .V., Rao, P. S., Ashan, M. M., Datta, R. K. and Rekha, M. (2000) Studies on combining ability and Reterosis in the silkworm, *Bombyx mori* L. *Indian J.Seric.*,**39**:43-48.

Singh Ravindra; Nagaraju. J., Ram Mohan Rao P., Premalatha ,V., Vijayaraghavan, K. and Sengupta, K. (1990) Heterosis analysis in the silkworm, *Bombyx mori* L. *Sericologia*,**30**: 293-300.

Singh, Ravindra; Vijayaraghavan, K., Premalatha, V., Ram Mohan Rao P., Sengupta, K. and Kannantha, V.(1992) Hybrid vigour in F1 , F2 and back crosses in silkworm, *Bombyx mori* L. *Mysore J. Agric.Sci.*,**26**:76- 81.

Singh Ravindra.; Raghavendra Rao, D., Premalatha, V., Mondal, Sipra, Karippa, B.K., Jayaswal, K. P. and Datta, R.K. (2001a) Manifestation of hybrid vigour and cocoon shape variability in F1hybrids of the mulberry silkworm, *Bombyx mori* L. *Int. J. Indust. Entomol.*,**2**:133-139.

Singh Ravindra.; Raghavendra Rao, D., Premalatha, V., Karippa, B.K., Jayaswal, K.P.and Datta, R.K. (2001b) Evaluation of combining ability in hybrids between low, medium and high cocoon weight polyvoltine and bivoltine breeds of silkworm, *Bombyx mori* L. *Sericologia*,**41**:57- 64.

Singh, Ravindra; Kalpana, G.V., Rao, P. S., Ashan, M. M., Datta, R. K. and Rekha, M. (2000) Studies on combining ability and Reterosis in the silkworm, *Bombyx mori* L. *Indian J.Seric.*,**39**:43-48.

Singh, Ravindra; Kalpana, G.V.; Rao,P.S., Ashan, M. M., Datta, R. K. and Rekha, M. (2000) Studies on combining ability and Reterosis in the silkworm, *Bombyx mori* L. *Indian J.Seric.*,**39**:43-48.1-37.

Sinha, A.K.and Choudhuri,A.(1992) Factors influencing the phenology of different broods of tropical tasar silk moth *Antheraea mylitta* Drury (*Lepidoptera:Saturniidae*) in relation to emergence and post emergence behaviour. *Environment and Ecology*,**10(4)**:952-958.

Srivastava,A. K.; Naqvi, A. H., Roy, G. C. and Sinha. B.R.R.P. (1988) Temporal variation in qualitative and quantitative characters of *Antheraea mylitta* Drury.The third international conference on wild silk moths,Vol 5,54-56

Srivastava, A. K.; Mishra, P. K., and Jayaswal, J.(1996) Life Table Studies of Muga Silkworm *A. assama* westwood (*Lepidoptera: Saturniidae*) on three different host plants,*Ann.Entomol.*,**14(1)** 3.

Subba Rao,G.and Sahai ,V.(1989) Combining ability and heterosis studies in bivoltine strains of silkworm *Bombyx mori* L. *Uttar Pradesh J. Zool.*

Tayade, D. S.(1989) Genetic architecture of economic traits in some strains of mulberry silkworm, *Bombyx mori* L. Combining ability analysis. *Sericologia*, **29**:43-60.

Tazima, Y. (1958) Report on *Sericulture Industry in India*, Central Silk Board, Bangalore, India, 23-26.

Thangavelu, K.(1991) Wild sericigenous insects of India: A need for, conservation. *Wild Silk moth*, 71-77.

Thangavelu, K.; Bajpayee, C.M.and Bania, H.R. (1992) Indoor rearing of tropical tassar silkworm, *A.mylitta*. D, *Wild silkmoths*, 99-103.

Thangavelu, K.:(1999), Silkworm Breeding : Priorities for India .In in Mulberry Sericulture. C.V.G. Publications., Bangalore, 187-207.

Tikoo, B.L.,Sitaram Iyengar, M.N.S., Dhar K.L. and Kaul, A.N. (1975) Seasonal effect of seed production on the hatchability of silkworm *Bombyx mori* (L). *Indian J. Seric.*,**14**:12-15.

Toyama, K. (1906)Breeding method of silkworm. *Sangyo-Shimpo*, 158:283-286.

Ueda S.; Kimura R.and Suzuki, k,. (1969) Studies on the growth of the silkworm, *Bombyx mori* , II. The influence of the rearing condition upon the larval growth, productivity of silk , substance and eggs and boiled off loss in the cocoon shell. *Bull. Sericult. Expt. Sta. Japan*, **23**:290-293.

Uvarov, B.P. (1931) Insects and climate. *Transactions of the Entomological Society, London*, **79**:1-247.

Virk, J.S. ; Rabinder Kaur and Parwinder Kaur. (2009) Performance of eri silk worm, *Samia cynthia ricini* Boisduval in différent seasons of Punjab. *Indian J.Seric.*, **48(1)**:78-80.

Watanabe K . Further studies on the voltinism in silkworm *Bombyx mori* (1928). Bull. Sericul. Expt. Sta . Japan, **7**:285-303.

Wigglesworth, V.B.(1973) *The principles of Insect Physiology*, 7th Edition chaman and Hall, London,Great Britain.*Indian.J.Seric.*,**1**:79-86.

Yamazaki (1939) A phonological study of non-mulberry silkworm. *J.Applied Zool.***11**:167-176.

Yokoyama,T. and Takashima, M.(1952) *Acta Sericologia* **2**:159-164.